



ENTAA Care

Ear, Nose & Throat, Asthma & Allergy
Audiology & Hearing Instrument Center
Speech & Balance Center

If you have any questions, regarding the effect of any medications on the list, please call our office @ 410-760-8840.

Please do not take any of the following **48 hours** before the test. However, **contact your physician should you have any reservations about discontinuing any medications.**

- | | |
|-------------------------------|---|
| Antibiotics: | Gentamicin, Streptomycin |
| Anticonvulsants: | Klonopin (do not stop taking for anxiety), Dilantin, Librium, Librax, Tegretol (carbamazepine), Nembutal |
| Antidepressants/Sedatives: | Ativan, Elavil, Lithium, Pamelor, Placidyl, Prozac, Valium, Xanax |
| Antihistamines/Decongestants: | Allegra, Benadryl, Chlor-Trimeton, Claritin, Seldane, Sudafed, Zyrtec, Xyzal, Over-the-Counter cold remedies |
| Anti-nausea medicine: | Dramamine, Compazine, Bonine, Marezine, Votrol, Phenergan, Thorazine, etc. |
| Anti-vertigo medicine: | Anti-vert, Meclizine, Ru-vert, Transderm Scop patches, Vontrol
Transderm Scop patches - must be off for 5 days prior to testing |
| Narcotics and Barbituates: | Phenobarbital, Codeine, Demerol, Dilaudid, Percocan, Percocet, Phenaphen, etc. |
| Sleeping Medications: | Ambien, Dalmane, Doral, Halcion, Lunesta, ProSom, Restoril, Seconal, Unisom |
| Stimulant Medications: | Amphetamine, Ritalin |
| Tranquilizers: | Valium, Librium, Atarax, Visteril, Equanil, Miltown, Travil, Serex, Etrafon, Ativan, etc. |
| Other Medications: | Alcohol (beer, wine, liquor, or cough medications containing alcohol)
Methadone, NSAID, Quinine, Salicilates-Aspirin |

No recreational drugs

**** Anti-Epilepsy and Anti-Psychotic medications, as well as medications for Diabetes, Hypertension, or Heart Disease should not be discontinued.**

ADMINISTRATIVE OFFICE

203 Hospital Drive, Suite 200 · Glen Burnie, MD 21061 · Phone (410) 760-8840 · Fax (410) 760-8847

www.entaacare.com

Annapolis

Columbia

Glen Burnie

Kent Island

Laurel

Odenton